

THE STARBURST CONTRIBUTION TO THE EXTRA-GALACTIC γ -RAY BACKGROUND

TODD A. THOMPSON,^{1,2} ELIOT QUATAERT,³ & ELI WAXMAN⁴

Draft version February 5, 2008

ABSTRACT

Cosmic ray protons interacting with gas at the mean density of the interstellar medium in starburst galaxies lose energy rapidly via inelastic collisions with ambient nuclei. The resulting pions produce secondary electrons and positrons, high-energy neutrinos, and γ -ray photons. We estimate the cumulative γ -ray emission from starburst galaxies. We find a total integrated background above 100 MeV of $F_\gamma \approx 10^{-6}$ GeV cm⁻² s⁻¹ sr⁻¹ and a corresponding specific intensity at GeV energies of $\nu I_\nu \approx 10^{-7}$ GeV cm⁻² s⁻¹ sr⁻¹. Starbursts may thus account for a significant fraction of the extra-galactic γ -ray background. We show that the FIR-radio correlation provides a strong constraint on the γ -ray emission from starburst galaxies because pions decay into both γ -rays and radio-emitting electron/positron pairs. We identify several nearby systems where the potential for observing γ -ray emission is the most favorable (M82, NGC 253, & IC 342), predict their fluxes, and predict a linear FIR- γ -ray correlation for the densest starbursts. If established, the FIR- γ -ray correlation would provide strong evidence for the “calorimeter” theory of the FIR-radio correlation and would imply that cosmic rays in starburst galaxies interact with gas at approximately the mean density of the interstellar medium (ISM), thereby providing an important constraint on the physics of the ISM in starbursts.

Subject headings: galaxies:starburst — gamma rays:theory, observations — cosmology:diffuse radiation — ISM:cosmic rays — radiation mechanisms:non-thermal

1. INTRODUCTION

The magnitude of the extra-galactic γ -ray background is uncertain, primarily due to the presence of foreground contaminants (compare Keshet et al. 2004, Strong et al. 2004, Sreekumar et al. 1998). *GLAST*⁵ will have an order of magnitude better sensitivity at GeV energies than previous experiments and should provide important constraints on the unresolved extra-galactic γ -ray emission. A number of potential contributors to the GeV γ -ray background have been discussed in the literature including blazars (Stecker & Salamon 1996), galaxy clusters and groups (Colafrancesco & Blasi 1998; Dar & Shaviv 1995), intergalactic shocks and structure formation (Loeb & Waxman 2000; Keshet et al. 2003; Miniati 2002), dark matter annihilation (e.g., Ullio et al. 2002; Elsässer & Mannheim 2005), and normal star-forming galaxies (Pavlidou & Fields 2002). In this paper we calculate the contribution of starburst galaxies to the GeV γ -ray background. In particular, we assess the possibility that starbursts are cosmic ray proton calorimeters: the dense ISM of these systems acts as a beam dump for the total energy injected by supernovae into cosmic ray protons, with a portion of the proton energy emerging as γ -rays and high-energy neutrinos.

In §2 we review Völk’s (1989) electron calorimeter model for the observed FIR-radio correlation (see Thompson et al. 2006, hereafter T06). We then argue that starbursts may also be proton calorimeters. In §3, we predict the γ -ray flux from starburst galaxies and highlight which nearby systems are most likely to have detectable γ -ray fluxes. We argue that

the observed FIR-radio correlation provides important constraints on the γ -ray emission from starbursts because pions from inelastic proton-proton collisions produce both γ -rays and secondary electrons and positrons, which then produce radio synchrotron. Section 4 then discusses the cumulative starburst contribution to the extra-galactic γ -ray background. Estimates similar to those presented here for the high-energy neutrino background from starbursts have recently been made by Loeb & Waxman (2006).

Although the γ -ray flux from individual starbursts has been estimated by several authors (e.g., Torres 2004, Torres et al. 2004, Cillis et al. 2005, Paglione et al. 1996, and Blom et al. 1999), this paper estimates the expected extra-galactic background, defines the requisite conditions for proton calorimetry, and explicitly connects the γ -ray predictions with constraints from the FIR-radio correlation.

2. STARBURST GALAXIES AS COSMIC RAY CALORIMETERS

Observations of both normal star-forming and starburst galaxies reveal a linear correlation between their radio (primarily synchrotron) and FIR luminosities (van der Kruit 1971, 1973; Helou et al. 1985; Condon 1992; Yun et al. 2001; Mur-
gia et al. 2005; Murphy et al. 2006). Völk (1989) argued that the FIR-radio correlation arises because the synchrotron cooling time for relativistic electrons generated in supernova shocks is significantly shorter than their escape time from the galactic disk (the “calorimeter theory”). In this limit, the cosmic ray electrons radiate nearly all of the energy supplied to them by supernovae. The calorimeter theory is a particularly compelling explanation for the FIR-radio correlation in starburst galaxies, where the synchrotron and inverse Compton (IC) cooling times of cosmic ray electrons are much shorter than in normal star-forming galaxies (T06).

A priori, one of the strongest objections to the calorimeter theory is that the radio spectra of galaxies are not compatible with the expected steepening due to strong synchrotron cooling (e.g., Condon 1992; Niklas & Beck 1997): in the presence of strong cooling, models predict $F_\nu \propto \nu^{-\alpha}$ with $\alpha \approx 1$,

¹ Department of Astrophysical Sciences, Peyton Hall-Ivy Lane, Princeton University, Princeton, NJ 08544; thomp@astro.princeton.edu

² Lyman Spitzer Jr. Fellow

³ Astronomy Department & Theoretical Astrophysics Center, 601 Campbell Hall, The University of California, Berkeley, CA 94720; eliot@astro.berkeley.edu

⁴ Physics Faculty, Weizmann Institute of Science, Rehovot 76100, Israel; waxman@wicc.weizmann.ac.il

⁵ <http://glast.gsfc.nasa.gov/>; <http://www-glast.stanford.edu/>

while observations find $\alpha \approx 0.75$, even in starbursts (e.g., Condon et al. 1991; Condon 1992; Niklas et al. 1997). T06 argue that if cosmic ray electrons interact with gas at approximately the mean density of the ISM, ionization and relativistic bremsstrahlung losses will systematically flatten the nonthermal spectra of starburst galaxies and simultaneously maintain the linearity of the FIR-radio correlation. This strongly supports Völk's calorimeter model for the FIR-radio correlation. In the model of T06, the conclusion that cosmic rays interact with gas at roughly the mean density of the ISM is required to account for the radio spectra of starbursts in the presence of strong synchrotron cooling. This requirement has an important consequence for cosmic ray protons, accelerated together with electrons in supernova remnants.

If cosmic ray protons interact with gas at the mean density of the ISM in starbursts, then they lose energy rapidly via inelastic proton-proton collisions with ambient nuclei. This interaction generates both charged and neutral pions, which subsequently produce secondary electrons and positrons, high energy neutrinos, and photons, respectively. For protons of GeV energy, the total inelastic proton-proton cross section is ≈ 30 mb (with inelasticity ~ 0.5 ; e.g., Gaisser 1990) so that the cosmic ray proton energy-loss timescale is

$$\tau_{pp} \approx 7 \times 10^7 n^{-1} \text{ yr}, \quad (1)$$

where n is the number density in units of cm^{-3} . The energy-loss timescale decreases by less than a factor of two over six orders of magnitude in proton energy.

If the pion production timescale within a galaxy is shorter than the escape timescale of relativistic protons, then the protons will lose most of their energy to secondary photons, electron/positron pairs, and neutrinos before escaping the galaxy. The galaxy would then be a cosmic ray proton calorimeter.

The escape timescale for cosmic ray protons is uncertain. Both energy-dependent diffusion and advection in a galactic outflow likely contribute. In the Galaxy, the diffusive escape timescale for 10 GeV protons is inferred to be $\tau_{\text{diff}} \sim 3 \times 10^7$ yr near the solar circle (Garcia-Munoz et al. 1977; Connell 1998). Such protons interact with a mean total column density of $\sim 10 \text{ g cm}^{-2}$ before escaping, and the column decreases at higher energies roughly as $E^{-1/2}$ (e.g., Engelmann et al. 1990; Webber et al. 2003). These observations imply that 10 GeV cosmic rays interact with matter of density $n \sim 0.2 \text{ cm}^{-3}$, somewhat below the average density of the ISM in the gas disk of the Milky Way. Using equation (1) we infer that 10 GeV protons lose roughly 10% of their energy to pion production before escaping. Assuming that diffusive losses proceed similarly in other star-forming galaxies, these results imply that there is a critical gas surface density Σ_{crit} above which protons will lose most of their energy to pion production. Taking the Milky Way's surface density to be $\Sigma_{\text{MW}\odot} \approx 2.5 \times 10^{-3} \text{ g cm}^{-2}$ at the solar circle (e.g., Boulares & Cox 1990) we estimate that

$$\Sigma_{\text{crit}}^{\text{diff}} \sim 10 \Sigma_{\text{MW}\odot} \approx 0.03 \text{ g cm}^{-2}. \quad (2)$$

For starburst galaxies, the assumption that diffusion dominates the escape of cosmic rays likely overestimates the escape timescale. In these systems escape is probably dominated by advection out of the gas mid-plane in a starburst-driven wind (as, e.g., in M82; see Klein et al. 1988, Seaquist & Odegard 1991). An estimate for the escape timescale is then

$$\tau_{\text{wind}} \approx h/V \approx 3 \times 10^5 h_{100} V_{300}^{-1} \text{ yr}, \quad (3)$$

where $h_{100} = h/100 \text{ pc}$ is the gas scale height and $V_{300} = V/300 \text{ km s}^{-1}$ is a typical wind velocity (e.g., Martin 1999). The condition $\tau_{pp} \lesssim \tau_{\text{wind}}$ is then the criterion for cosmic ray protons to lose a significant fraction of their energy to pion production. This criterion may also be written in terms of a critical gas surface density;

$$\Sigma_{\text{crit}}^{\text{wind}} \approx 0.3 V_{300} \text{ g cm}^{-2}. \quad (4)$$

Although uncertain, equations (2) and (4) imply that galaxies with $\Sigma_g \gtrsim 0.03 - 0.3 \text{ g cm}^{-2}$ are likely to be proton calorimeters. A number of local starburst galaxies satisfy this criterion (see Table 1).

3. γ -RAY EMISSION FROM STARBURST GALAXIES & THE FIR- γ -RAY CORRELATION

To estimate the γ -ray emission associated with pion production in starbursts, we assume that the star formation rate ($\dot{M}_*$) is related to the total IR luminosity ($L_{\text{TIR}} [8 - 1000] \mu\text{m}$; see, e.g., Calzetti et al. 2000, Dale et al. 2001) by $L_{\text{TIR}} = \epsilon \dot{M}_* c^2$ (ϵ is an IMF-dependent constant), that the supernova rate per unit star formation, Γ_{SN} , is a constant fraction of $\dot{M}_*$, and that $5\eta_{0.05}\%$ of the energy of each supernova ($E_{51} = E_{\text{SN}}/10^{51} \text{ ergs}$) is supplied to relativistic protons. With these assumptions, the total energy per unit time injected in cosmic ray protons is

$$L_{\text{CR}_p} \approx 4.6 \times 10^{-4} L_{\text{TIR}} E_{51} \eta_{0.05} \beta_{17}, \quad (5)$$

where $\beta_{17} = (\Gamma_{\text{SN}}/\epsilon)/17 M_{\odot}^{-1}$; for continuous star formation over $\sim 10^8$ yrs, this ratio has only a very weak dependence on the assumed IMF, since high mass stars dominate both the total luminosity and the supernova rate. In addition, for star-formation timescales of $3 \times 10^7 - 10^9$ yrs, β increases by only a factor of approximately 1.5, from ≈ 15 to ≈ 23 (e.g., Leitherer et al. 1999). We further assume that starburst galaxies are proton calorimeters ($\tau_{pp} < \tau_{\text{esc}}$, $\Sigma_g > \Sigma_{\text{crit}}$) and — for the purposes of an analytic estimate — that the γ -ray spectrum is flat at and above GeV energies; this requires that cosmic ray protons are injected with a spectral index $p \approx 2$ (e.g., Blandford & Eichler 1987). Numerical calculations for different p are shown in Figure 1 and discussed below.

Because the cross-sections for neutral- and charged-pion production are similar, $\approx 1/3$ of the proton energy goes into neutral pions, which subsequently decay into γ -rays. Thus, under the assumption of proton calorimetry, the expected γ -ray luminosity can be directly related to L_{CR_p} and L_{TIR} (eq. [5]). For a flat spectrum and for energies $\gtrsim \text{GeV}$, we find that

$$\nu L_{\nu}(\text{GeV}) \approx 1.5 \times 10^{-4} \frac{\eta_{0.05} L_{\text{TIR}}}{\ln(\gamma_{\text{max}})} \approx 10^{-5} \eta_{0.05} L_{\text{TIR}}, \quad (6)$$

where $\gamma_{\text{max}} (= 10^6)$ is the assumed maximum Lorentz factor of the accelerated protons and we have dropped the dependence here and below on the ratio $\beta = (\Gamma_{\text{SN}}/\epsilon)$.

Approximately 2/3 of the energy in cosmic ray protons goes to charged pions, which subsequently decay into high energy neutrinos and electron/positron pairs. The pairs receive $\approx 1/4$ of the pion energy (Schlickeiser 2002). Thus $\approx 0.8\eta_{0.05}\%$ of the energy of each supernova goes into secondary electrons and positrons. Assuming that the synchrotron cooling time of the secondary pairs is less than their escape time (Völk 1989; T06), we find a FIR-radio correlation of the form

$$\nu L_{\nu}(\text{GHz}) \approx 7.5 \times 10^{-5} \frac{\eta_{0.05} L_{\text{TIR}}}{2 \ln(\gamma_{\text{max}})} \approx 2.5 \times 10^{-6} \eta_{0.05} L_{\text{TIR}}. \quad (7)$$

The observed correlation is $\nu L_\nu(1.4 \text{ GHz}) \approx 1.1 \times 10^{-6} L_{\text{TIR}}$ (Yun et al. 2001).⁶ For $\eta = 0.05$, equation (7) somewhat over-predicts the observed FIR-radio correlation in starbursts. However, this prediction neglects ionization, inverse Compton, and bremsstrahlung losses, which may be competitive with synchrotron losses in starbursts and will reduce the magnitude of $\nu L_\nu(\text{GHz})/L_{\text{TIR}}$.⁷ For example, in the models of T06, which simultaneously explain the linearity of the FIR-radio correlation and the radio spectra of starbursts (their Figs. 3 & 4), bremsstrahlung and ionization losses account for $\approx 50\%$ of the energy lost by cosmic ray electrons. Note that even if inverse Compton and bremsstrahlung losses completely dominate ionization and synchrotron losses for secondary electrons/positrons, their contribution to the total X-ray and γ -ray emission from starbursts is at most 1/2 of the contribution from neutral-pion decay; inverse Compton and bremsstrahlung will, however, dominate the emission below $\sim 100 \text{ MeV}$.

Equations (6) and (7) highlight the importance of the FIR-radio correlation for constraining the γ -ray emission from starbursts. Statistically, the γ -ray emission from galaxies cannot exceed that predicted by equation (6) without violating the radio constraints implied by the FIR-radio correlation (eq. 7). The only way out of this conclusion is if ionization, inverse Compton, and bremsstrahlung losses significantly exceed synchrotron losses for cosmic ray electrons and positrons. Although possible, it would be remarkable if synchrotron losses were highly sub-dominant in starbursts and yet somehow the linearity of the FIR-radio correlation is maintained, both within different starburst galaxies and between starbursts and normal star-forming galaxies.

Equation (7) is only consistent with the observed FIR-radio correlation for $\eta \sim 0.05$, i.e., if $\sim 5\%$ of the supernova energy is supplied to relativistic protons. Although this efficiency of cosmic ray production is reasonably consistent with the energetics of galactic cosmic rays (e.g., Schlickeiser 2002) and with the γ -ray luminosity of the Milky Way (Hunter et al. 1987), some models of non-linear diffusive shock acceleration imply $\eta \sim 0.5$ (e.g., Ellison & Eichler 1984; Ellison et al. 2004).⁸ If indeed $\eta \gg 0.05$, then the FIR-radio correlation implies that either starburst galaxies are not proton calorimeters or that synchrotron losses are highly sub-dominant with respect to ionization, inverse Compton, and bremsstrahlung losses.

Equations (6) and (7) imply that the γ -ray fluxes of the densest starbursts should be linearly proportional to their radio and IR fluxes in the proton calorimeter limit. Table 1 provides a sample of galaxies chosen to have large IR fluxes and gas densities (see also Paglione et al. 1996; Blom et al. 1999; Torres et al. 2004; Torres 2004; Domingo-Santamaría & Torres 2005; Torres & Domingo-Santamaría 2005). For the latter, we have applied a cut $\Sigma_g \geq 0.02 \text{ g cm}^{-2}$, plausibly in the proton calorimeter limit; the implied mean

gas densities range from $\sim 20 \text{ cm}^{-3}$ for IC 342 to $\sim 10^4 \text{ cm}^{-3}$ for Arp 220. The systems in Table 1 are listed in order of decreasing TIR flux density, which corresponds to decreasing γ -ray flux in the proton calorimeter limit (eq. [6]). Several systems just miss our cut on Σ_g , including the nucleus of NGC 5236, NGC 4631, and NGC 3521.⁹ Because typical errors in Σ_g are of order $\pm 0.2 \text{ dex}$ (e.g., Kennicutt 1998) and because of uncertainty in Σ_{crit} (§2), these systems may also prove to be bright sources in the γ -ray sky.

The γ -ray fluxes predicted by equation (6) at 1 GeV from L_{TIR} are also provided in Table 1. For comparison, the expected sensitivity of *GLAST* at 1 GeV for a 1 year integration is $\approx 2 \times 10^{-10} \text{ GeV cm}^{-2} \text{ s}^{-1}$ (see Fig. 1).¹⁰ Table 1 shows that there are 3 nearby starbursts above this limit (M82, NGC 253, & IC 342) and approximately 10 other systems with predicted fluxes within a factor of 4 of *GLAST*'s detection limit. We note that NGC 3690 and Arp 220 would be particularly important for testing our predicted FIR- γ -ray correlation over a large dynamic range in L_{TIR} .

Figure 1 shows the γ -ray spectrum, as calculated using the formalism of Aharonian & Atoyan (2000) (their §4.1), as a function of energy for the specific cases of NGC 253 (*dashed lines*) and Arp 220 (*dotted lines*), for $p = 2.0 - 2.4$, where p is the slope of the proton injection spectrum. We first specify p , and then we calculate the spectrum from equation (22) of Aharonian & Atoyan (2000), adopting their fit for the cross section and their value of $K_\pi \approx 0.17$ (see also the Appendix of Domingo-Santamaría & Torres 2005). We normalize the resulting spectrum, such that the integral over all γ -ray energies is set by the second equality in equation (6). Comparing our analytic estimate in equation (6) with this numerical calculation, we find that a nearly flat $p = 2.1$ proton spectrum gives the best agreement at GeV energies.

In order to be consistent with the FIR-radio correlation as given in the second equality in equation (7), all calculations with $p \neq 2.1$ are normalized to the $p = 2.1$ spectrum at 1 GeV. Note that this constraint implies that p and η are *not independent*. A constant flux at 1 GeV requires $(\eta, p) \approx (0.09, 2.0), (0.05, 2.1), (0.03, 2.2), (0.025, 2.3)$, and $(0.02, 2.4)$. Because the average magnetic field strengths in starbursts may range from $B \sim 0.1 \text{ mG}$ (e.g., NGC 253 or M82) to $B \sim 10 \text{ mG}$ (e.g., Arp 220) (see T06 for a discussion), and because the energy of GHz-emitting electrons/positrons is proportional to $B^{-1/2}$, our normalization of the $p \neq 2.1$ curves at 1 GeV (rather than the proper normalization of the 1 GHz radio emission) introduces a factor of ≈ 2 uncertainty in the predicted γ -ray emission.

As Figure 1 emphasizes, NGC 253 (and M82, see Table 1) should be readily detected with *GLAST*. Existing observations by *HESS* at TeV energies (Aharonian et al. 2005) yield an upper limit of $< 1.9 \times 10^{-12} \text{ ph cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$ (for a point source; $E_\gamma > 300 \text{ GeV}$) for NGC 253 that is very close to our prediction for $p = 2$, suggesting that longer integrations with *HESS* should be able to detect NGC 253 or significantly constrain the high energy spectral slope (see also Domingo-Santamaría & Torres 2005; Torres & Domingo-Santamaría 2005).

In two cases — NGC 253 and Arp 220 — we can compare

⁶ L_{FIR} , as defined in Yun et al. (2001) from the $60\mu\text{m}$ and $100\mu\text{m}$ IRAS fluxes (Helou et al. 1988), is generally a factor of ≈ 1.75 smaller than L_{TIR} for starburst galaxies (see Calzetti et al. 2000).

⁷ Equation (7) also neglects the contribution to the radio emission from primary electrons accelerated directly in supernova shocks. Typical estimates for the ratio of the total energy injected into protons and primary electrons are $\sim 20 - 40$ (e.g., Beck & Krause 2005), in which case secondary electrons/positrons likely dominate the radio emission in starbursts (see T06; Rengarajan 2005).

⁸ However, Völk et al. (2003) argue that standard spherical models of non-linear shock acceleration over-estimate the efficiency of cosmic ray production by a factor of ~ 5 .

⁹ The Seyfert 2 galaxy NGC 1068 with $\Sigma_g \approx 0.1 \text{ g cm}^{-2}$ (Helfer et al. 2003) would be fourth in Table 1 with $\nu F_\nu(\text{GeV}) \approx 11.8 f \times 10^{-11} \text{ GeV cm}^{-2} \text{ s}^{-1}$ where f is the fraction of the infrared emission due to the nuclear starburst rather than the AGN.

¹⁰ See <http://people.roma2.infn.it/~glast/> for a comparison of γ -ray telescope sensitivities.




FIG. 1.— γ -ray flux and specific intensity as a function of γ -ray energy. The data points with error bars show the extra-galactic γ -ray background as inferred by Strong et al. (2004) from the *EGRET* data. The open square shows the upper limit on the background derived by Keshet et al. (2004) at 1 GeV. The thick solid lines are the point source sensitivities of *EGRET* and *GLAST* for a one year all-sky survey and the *HESS* sensitivity curve for a 5σ point source detection using a 50 hour integration. The solid lines show the total starburst background calculated with $p = 2.0 - 2.4$ and assuming $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $f(z) = \min[1, 0.9z + 0.1]$, and $\dot{\rho}_*(z)$ from RSF2 in Porciani & Madau (2001). For this model $f_{\text{cal}} \approx 0.8$ (see eq. 10). The expected signal from the starbursts NGC 253 (dashed lines) and Arp 220 (dotted lines) with $p = 2.0 - 2.4$ (see §2, eqs. [6] & [7]) are also shown. For other starburst galaxies potentially visible by *GLAST* and *HESS*, see Table 1. For both the γ -ray background and the individual starbursts, the $p = 2.1$ curves are normalized such that the total integrated γ -ray luminosity is $1.5 \times 10^{-4} \eta_{0.05}$ times the total integrated stellar luminosity (eq. [6]). Because of the constraints on the radio emission from starbursts, all $p \neq 2.1$ curves are normalized to the $p = 2.1$ calculation at 1 GeV, which implies that in our models, p and η are not independent (see discussion in §2). Note that the effects of attenuation of the background spectrum above ~ 10 GeV have been neglected (see, e.g., Madau & Phinney 1996).

our estimate for the γ -ray luminosity with more detailed calculations by Torres (2004) and Domingo-Santamaría & Torres (2005). To make an apposite comparison, it is necessary to scale our results to their assumed parameters. Using $\eta = 0.165$ and $p = 2.2$, Torres (2004) finds for Arp 220 that $\nu F_\nu(\text{GeV}) \approx 3 \times 10^{-10} \text{ GeV cm}^{-2} \text{ s}^{-1}$. Rescaling our $\eta \approx 0.03$ and $p = 2.2$ result, and adjusting for our somewhat ($\approx 20\%$) higher TIR flux, we find that $\nu F_\nu(\text{GeV}) \approx 2.8 \times 10^{-10} \text{ GeV cm}^{-2} \text{ s}^{-1}$. For NGC 253, Domingo-Santamaría & Torres (2005) estimate $\nu F_\nu(\text{GeV}) \approx 2 \times 10^{-9} \text{ GeV cm}^{-2} \text{ s}^{-1}$ with $\eta = 0.1$ and $p = 2.2$. Again scaling our results, we find that $\nu F_\nu(\text{GeV}) \approx 2.3 \times 10^{-9} \text{ GeV cm}^{-2} \text{ s}^{-1}$ for their assumed parameters. The correspondence between our predicted fluxes and these more detailed calculations is thus excellent. These studies also find, in agreement with our assumptions, that π^0 decay dominates γ -ray emission at GeV energies, that secondary electrons and positrons dominate the radio emission at GHz frequencies, and that both systems are safely in the proton and

electron/positron calorimeter limits; the cooling timescale is less than the escape timescale.

The predicted fluxes for Arp 220 and NGC 253 above – for Torres (2004) and Domingo-Santamaría & Torres’s (2005) parameters – are a factor of ≈ 5 and 3 times higher than those given in Table 1 (primarily because of the larger values of η). Given the constraints imposed by the FIR-radio correlation, it is therefore perhaps surprising that Torres (2004) and Domingo-Santamaría & Torres’s (2005) are able to fit the radio emission from Arp 220 and NGC 253 with the significantly larger values of η they assumed. As per the discussion after equation (7), this is only possible in the proton calorimeter limit if ionization, bremsstrahlung, and/or inverse Compton losses dominate synchrotron losses. Indeed, for the individual nuclei of Arp 220 and for the inner starburst of NGC 253, Torres (2004) and Domingo-Santamaría & Torres’s (2005) parameters (gas densities, magnetic field strengths, etc.) imply that the ratio of the synchrotron cooling timescale

to the total electron energy loss timescale (including all processes; dominated by ionization losses) is ~ 5 . Thus, their results are fully consistent with the arguments of our paper. However, we believe that their predicted γ -ray fluxes are optimistic because, as noted above, the tightness and linearity of the FIR-radio correlation are difficult to explain if synchrotron losses are highly sub-dominant in all star-forming galaxies.

4. THE STARBURST γ -RAY BACKGROUND

The total diffuse extra-galactic TIR background from star formation is typically estimated to be $F_{20}^{\text{TIR}} = F^{\text{TIR}}/20 \text{ nW m}^{-2} \text{ sr}^{-1} \approx 1$, with roughly half of the contribution to the background coming from $z > 1$ (e.g., Nagamine et al. 2006). Equation (6) for the γ -ray flux from a given starburst in the proton calorimeter limit then implies a total integrated γ -ray background of

$$F_\gamma \approx 1.4 \times 10^{-6} \eta_{0.05} f_{0.75} F_{20}^{\text{TIR}} \text{ GeV s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1} \quad (8)$$

where $f_{0.75} = f_{\text{cal}}/0.75$ and

$$f_{\text{cal}} = \int_0^\infty \frac{\dot{\rho}_*(z) f(z) dz}{[(1+z)^2 E(z)]} / \left\{ \int_0^\infty \frac{\dot{\rho}_*(z) dz}{[(1+z)^2 E(z)]} \right\} \quad (9)$$

is the fraction of the TIR extragalactic background produced by starburst galaxies that are in the proton calorimeter limit (i.e., $\Sigma_g > \Sigma_{\text{crit}}$; §2), $f(z)$ is that fraction at every z , $\dot{\rho}_*$ is the comoving star formation rate density, and $E(z) = [\Omega_m(1+z)^3 + \Omega_\Lambda]^{1/2}$. For a flat γ -ray spectrum, equation (8) implies a specific intensity at GeV energies of

$$\nu I_\nu(\text{GeV}) \approx 10^{-7} \eta_{0.05} f_{0.75} F_{20}^{\text{TIR}} \text{ GeV s}^{-1} \text{ cm}^{-2} \text{ sr}^{-1}. \quad (10)$$

The estimated γ -ray background depends sensitively on the fraction of star formation that occurs in the proton calorimeter limit, i.e., on $f(z)$ and f_{cal} . We estimate $f(z=0) \sim 0.1$ from the fraction of the local FIR and radio luminosity density produced by starbursts (Yun et al. 2001). Confirmation that local starbursts like M82 and NGC 253 are γ -ray emitters at the level predicted in Table 1 would significantly solidify the background predictions of equations (8) and (10) by confirming that such starbursts are indeed proton calorimeters. At high redshift, a much larger fraction of star formation occurs in high surface density systems likely to be calorimeters. For example, the strong evolution of the IR luminosity function with redshift indicates that F^{TIR} is dominated by luminous infrared galaxies at $z \sim 1$ (Dole et al. 2006); these systems are typical of those we expect to be proton calorimeters, in which case $f(z) \sim 1$ at $z \sim 1$. In addition, the typical gas surface density of UV selected galaxies, which account for a substantial fraction of the star formation at high redshift, is roughly $\Sigma_g \sim 0.06 M_{10.5} R_{6\text{kpc}}^{-2} \text{ g cm}^{-2}$, where $M_{10.5}$ is the inferred average gas mass in units of $10^{10.5} M_\odot$ and $R_{6\text{kpc}} = R/6 \text{ kpc}$ is the half-light radius (Erb et al. 2006ab). This surface density is comparable to Σ_{crit} estimated in §2. Thus, if we take $f(z \gtrsim 1) \sim 1$ and $\dot{\rho}_*(z)$ as in, e.g., Porciani & Madau (2001), we find that $f_{\text{cal}} \sim 1$ (eq. [10]), with a significant contribution to the γ -ray background coming from $z \approx 1-2$, so that

$$F_\gamma \propto \nu I_\nu(\text{GeV}) \propto f(z \approx 1-2). \quad (11)$$

More specifically, taking $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$, and assuming a function $f(z) = \min(0.9z + 0.1, 1)$, which smoothly interpolates from a small local starburst fraction ($f(z \approx 0) \approx 0.1$) to an order unity starburst fraction at high redshift, we find that $f_{\text{cal}} \approx 0.8$ for models RSF1, RSF2, and RSF3 from Porciani & Madau (2001) and for the “Fossil” model of Nagamine et

al. (2006). These models also all have $F_{20}^{\text{TIR}} \approx 1$. As an alternative, assuming $f(z) = \min[0.1(1+z)^3, 0.8]$, in which the starburst fraction increases roughly in proportion to $\dot{\rho}_*(z)$, $f_{\text{cal}} \approx 0.6$ for the same models.¹¹

Using a different methodology, Pavlidou & Fields (2002) calculate the contribution to the γ -ray background from star-forming galaxies and find a value for νI_ν at GeV energies that is 4–5 times higher than that given in equation (10). Pavlidou & Fields (2002) did not account for the constraints imposed by the FIR-radio correlation, nor do they explicitly discuss the importance of proton calorimetry and the essential competition between escape and pion losses (§2). By analogy with the Galaxy, Pavlidou & Fields (2002) assume that few GeV protons lose roughly 10% of their energy before escaping in all star-forming galaxies in the local universe ($z \approx 0$). Because roughly 10% of the local TIR luminosity density is generated by starburst galaxies, and because we assume that all starbursts are calorimeters, local starburst and normal star-forming galaxies contribute roughly equally to the γ -ray background at $z \approx 0$. As we have argued above, however, starbursts are the dominant mode of star formation at high z and thus $z \sim 1-2$ starbursts dominate the contribution of star-forming galaxies to the γ -ray background.

5. DISCUSSION

Figure 1 shows the numerically calculated γ -ray background as a function of γ -ray energy (*solid lines*), for $p = 2.0-2.4$ (see caption for details). The inferred background from *EGRET* from Strong et al. (2004), as well as the conflicting upper limit at 1 GeV derived by Keshet et al. (2004) are also shown.

Our results indicate that γ -ray production from pion decay in starburst galaxies contributes significantly to the observed background above 100 MeV. The considerable uncertainties in the γ -ray background determination, primarily because of foreground subtraction (see Keshet et al. 2004), complicate a more direct comparison. An important point of this paper is that the absolute starburst γ -ray flux — both that from individual galaxies and that of the background — cannot exceed that predicted in Figure 1 and Table 1 without over-producing the radio emission from secondary electrons and positrons produced in charged pion decay (see §3, eq. [7]). This conclusion is independent of whether starbursts are in fact proton calorimeters (as we have assumed), and can only be circumvented if ionization, inverse Compton, and bremsstrahlung losses significantly exceed synchrotron losses for cosmic ray electrons and positrons in starbursts; such a determination would put strong constraints on the origin of the FIR-radio correlation (T06; §3). If detected by *GLAST*, the low energy γ -ray emission from systems like NGC 253 and M82 (§2,

¹¹ Stecker (2006) presents upper limits on the γ -ray and neutrino backgrounds from starbursts that are significantly lower than our estimates here (see §5 for our neutrino background estimate). There are two reasons for this difference. First, Stecker advocates that 22% of the extragalactic TIR background is produced at $z > 1.2$ (see his Table 1) whereas we find that all models for $\dot{\rho}_*(z)$ that are consistent with observations of star formation at high- z and the local stellar mass density imply that roughly half of the TIR background is produced at $z > 1$ (e.g., RSF1, RSF2, and RSF3 from Porciani & Madau (2001), the “Fossil” model from Nagamine et al. 2006, the dust-corrected star formation history given by Schiminovich et al. 2005, and the analytic model given by Hernquist & Springel 2003). Second, we argue that a fraction of order unity of all star formation at $z \approx 1$ occurs in starbursts likely to be calorimeters (our $f(z)$; eq. 11), whereas Stecker quotes 13% in the range $0.2 \leq z \leq 1.2$ and 60% for $z > 1.2$ (his Table 1). His step function model for $f(z)$ significantly underestimates the contribution of $z \sim 1$ galaxies to the γ -ray background.

Fig. 1, Table 1), may help constrain the contribution of inverse Compton and bremsstrahlung to electron/positron losses.

The primary physical uncertainty in our estimate of the γ -ray fluxes from starbursts lies in whether cosmic rays do in fact interact with gas at approximately the mean density of the ISM. In particular, given that galactic winds efficiently remove mass and metals from galaxies (e.g., Heckman et al. 1990), it is unclear whether the cosmic rays actually interact with the bulk of the ISM, which is required for pion production to be significant (§2). Detection of γ -rays from starbursts at the level predicted in this paper would thus provide an important constraint on the physics of the ISM in starbursts and on the coupling between supernovae and the dense ISM, which contains most of the mass. In addition, the ionization and bremsstrahlung energy-loss timescales for electrons/positrons are similar to the proton-proton pion production timescale. Thus, constraints on pion production in starbursts via γ -ray observations directly constrain the importance of ionization and bremsstrahlung losses for shaping the radio emission from starburst galaxies (T06). By contrast, detection of (or upper limits on) the γ -ray emission from starbursts well below our predictions would rule out the hypothesis that starbursts are proton calorimeters, although they would, in our opinion, still be electron calorimeters. The alternative possibility suggested by equation (6), that $\eta \ll 0.05$, is unlikely, given the energetics of Galactic cosmic ray production (e.g., Strong et al. 2004).

We note that the total IR background from the particular model of the star formation history of the universe used to produce the γ -ray background in Figure 1 is $F^{\text{TIR}} \approx 2 \times 10^{-5}$ ergs cm $^{-2}$ s $^{-1}$ sr $^{-1}$, or ≈ 20 nW m $^{-2}$ sr $^{-1}$, with approximately 80% ($f_{\text{cal}} \approx 0.8$) coming from starburst galaxies. A complete calculation, accounting for the contribution from the old stellar population (which contributes significantly to the back-

ground at $z \approx 0$), finds a total background in starlight that is roughly twice as large (e.g., Nagamine et al. 2006). Equation (7) and, indeed, the existence of the FIR-radio correlation, implies that the expected radio background is $\nu L_{\nu}(\text{radio}) \approx 7.5 \times 10^{-10} \eta_{0.05} / [2 \ln(\gamma_{\text{max}})] \approx 2.7 \times 10^{-11}$ ergs cm $^{-2}$ s $^{-1}$ sr $^{-1}$. A factor of two reduction in $\nu L_{\nu}(\text{radio})$ has been included in this estimate as per the discussion after equation (7).

In addition to producing secondary electron/positron pairs and γ -rays, pion production is also a significant source of high-energy neutrinos, with $\approx 5\%$ of the proton cosmic ray energy going into neutrinos. Two-thirds of this energy goes to muon-type neutrinos. Therefore, for starburst galaxies that are proton calorimeters, we expect a FIR-neutrino correlation of the form $\nu L_{\nu}(\mu - \text{neutrinos}) = (2/3) \nu L_{\nu}(\text{all neutrinos}) \approx 1.5 \times 10^{-4} \eta_{0.05} L_{\text{TIR}} / \ln(\gamma_{\text{max}}) \approx 10^{-5} \eta_{0.05} L_{\text{TIR}}$, identical to the γ -ray luminosity (eq. [6]). The corresponding cumulative extra-galactic μ -neutrino background at GeV energies is then the same as given in equation (10). Using a very similar estimate, Loeb & Waxman (2006) have recently argued that starbursts are likely to be an important contributor to the high-energy neutrino background. Because the ratio of the γ -ray to neutrino luminosity from pion decay depends only on the micro-physics of pion production, the γ -ray fluxes from starbursts predicted in this paper will provide a crucial calibration of the expected flux of extra-galactic high energy neutrinos.

T.A.T. thanks Uri Keshet for a critical reading of the text and stimulating conversation. E.Q. is supported in part by NASA grant NNG06GI68G, an Alfred P. Sloan Fellowship, and the David and Lucile Packard Foundation. E.W. thanks the Institute for Advanced Study and is supported in part by ISF and AEC grants. We thank the anonymous referee for a prompt and useful report.

REFERENCES

- Aharonian, F. A., & Atoyan, A. M. 2000, *A&A*, 362, 937
 Aharonian, F., et al. 2005, *A&A*, 437, L7
 Beck, R., & Krause, M. 2005, *Astronomical Notes*, 326, 414
 Bell, E. F. 2003, *ApJ*, 586, 794
 Boulares, A., & Cox, D. P. 1990, *ApJ*, 365, 544
 Blandford, R., & Eichler, D. 1987, *Phys. Rep.*, 154, 1
 Blom, J. J., Paglione, T. A. D., & Carramiñana, A. 1999, *ApJ*, 516, 744
 Calzetti, D., Armus, L., Bohlin, R. C., Kinney, A. L., Koornneef, J., & Storchi-Bergmann, T. 2000, *ApJ*, 533, 682
 Cillis, A. N., Torres, D. F., & Reimer, O. 2005, *ApJ*, 621, 139
 Colafrancesco, S., & Blasi, P. 1998, *Astroparticle Physics*, 9, 227
 Condon, J. J. 1987, *ApJS*, 65, 485
 Condon, J. J., & Broderick, J. J. 1988, *AJ*, 96, 30
 Condon, J. J., Helou, G., Sanders, D. B., & Soifer, B. T. 1990, *ApJS*, 73, 359
 Condon, J. J., Huang, Z.-P., Yin, Q. F., & Thuan, T. X. 1991, *ApJ*, 378, 65
 Condon, J. J. 1992, *ARA&A*, 30, 575
 Condon, J. J., Helou, G., Sanders, D. B., & Soifer, B. T. 1996, *ApJS*, 103, 81
 Condon, J. J., et al. 1998, *AJ*, 115, 1693
 Connell, J. J. 1998, *ApJ*, 501, L59
 Dar, A., & Shaviv, N. J. 1995, *Physical Review Letters*, 75, 3052
 Dole, H., et al. 2006, *A&A*, 451, 417
 Domingo-Santamaría, E., & Torres, D. F. 2005, *A&A*, 444, 403
 Ellison, D. C., Decourchelle, A., & Ballet, J. 2004, *A&A*, 413, 189
 Ellison, D. C., & Eichler, D. 1984, *ApJ*, 286, 691
 Elsässer, D., & Mannheim, K. 2005, *Physical Review Letters*, 94, 171302
 Gaisser, T. K. 1990, *Cosmic Rays and Particle Physics*, (Cambridge: Cambridge University Press)
 García-Munoz, M., Mason, G. M., & Simpson, J. A. 1977, *ApJ*, 217, 859
 Golla, G., & Wielebinski, R. 1994, *A&A*, 286, 733
 Heckman, T. M., Armus, L., & Miley, G. K. 1990, *ApJS*, 74, 833
 Helfer, T. T., et al. 2003, *ApJS*, 145, 259
 Helou, G., Soifer, B. T., & Rowan-Robinson, M. 1985, *ApJ*, 298, L7
 Helou, G., Khan, I. R., Malek, L., & Boehmer, L. 1988, *ApJS*, 68, 151
 Hernquist, L., & Springel, V. 2003, *MNRAS*, 341, 1253
 Hunter, S. D., et al. 1997, *ApJ*, 481, 205
 Kennicutt, R. C. 1998, *ApJ*, 498, 541
 Keshet, U., Waxman, E., Loeb, A., Springel, V., & Hernquist, L. 2003, *ApJ*, 585, 128
 Keshet, U., Waxman, E., & Loeb, A. 2004, *JCAP*, 4, 6
 Klein, U., Wielebinski, R., & Morsi, H. W. 1988, *A&A*, 190, 41
 Leitherer, C., et al. 1999, *ApJS*, 123, 3
 Loeb, A., & Waxman, E. 2000, *Nature*, 405, 156
 Loeb, A., & Waxman, E. 2006, *JCAP*, 5, 3
 Madau, P., & Phinney, E. S. 1996, *ApJ*, 456, 124
 Martin, C. L. 2005, *ApJ*, 621, 227
 Miniati, F. 2002, *MNRAS*, 337, 199
 Murgia, M., Helfer, T. T., Ekers, R., Blitz, L., Moscadelli, L., Wong, T., & Paladino, R. 2005, *A&A*, 437, 389
 Murphy, E. J., et al. 2006, *ApJ*, 638, 157
 Nagamine, K., Ostriker, J. P., Fukugita, M., & Cen, R. 2006, *arXiv:astro-ph/0603257*
 Niklas, S., & Beck, R. 1997, *A&A*, 320, 54
 Paglione, T. A. D., Marscher, A. P., Jackson, J. M., & Bertsch, D. L. 1996, *ApJ*, 460, 295
 Pavlidou, V., & Fields, B. D. 2002, *ApJ*, 575, L5
 Porciani, C., & Madau, P. 2001, *ApJ*, 548, 522
 Schiminovich, D., et al. 2005, *ApJ*, 619, L47
 Seaquist, E. R., & Odegard, N. 1991, *ApJ*, 369, 320
 Sreekumar, P., et al. 1998, *ApJ*, 494, 523
 Stecker, F. W., & Salamon, M. H. 1996, *ApJ*, 464, 600
 Stecker, F. 2006, *arXiv:astro-ph/0607197*
 Strong, A. W., Moskalenko, I. V., & Reimer, O. 2000, *ApJ*, 537, 763
 Strong, A. W., Moskalenko, I. V., & Reimer, O. 2004, *ApJ*, 613, 956
 Thompson, T. A., Quataert, E., Waxman, E., Murray, N., & Martin, C. L. 2006, *ApJ*, 645, 186
 Torres, D. F., Reimer, O., Domingo-Santamaría, E., & Digel, S. W. 2004, *ApJ*, 607, L99
 Torres, D. F., 2004, *ApJ*, 617, 966

TABLE 1
PROPERTIES OF LOCAL STARBURST & STAR-FORMING GALAXIES

Object Name	D^a (Mpc)	Σ_g^b (g cm^{-2})	$S_{60\mu\text{m}}^c$ (Jy)	$S_{100\mu\text{m}}^d$ (Jy)	S_{TIR}^e (Jy)	$S_{1.4\text{GHz}}^f$ (mJy)	$\nu F_\nu(\text{GeV})^g$ ($10^{-11} \text{ GeV cm}^{-2} \text{ s}^{-1}$)	Data Refs.
NGC 3034 (M82)...	3.3	0.69	1313.5	1355.4	8302.4	7657.0	73	4,2,3
NGC 253.....	3.5	0.47	997.7	1857.8	7755.8	5594.0	68 ^h	1,2,3
IC 342.....	4.4	0.02	255.9	660.7	2311.6	2250.0	20	1,2,3
NGC 2146.....	12.6	0.14	144.5	204.2	1009.8	1074.6	8.9	1,2,3
NGC 3690.....	42.2	0.04	119.7	118.6	748.0	678.3	6.6	4,2,3
NGC 1808.....	14.2	0.09	104.5	147.2	729.4	529.0	6.4	1,2,3
NGC 1365.....	18.6	0.09	84.1	185.4	704.2	530.0	6.2	7,8,3
Arp 220.....	76.6	12.0	107.4	118.3	691.9	326.9	6.1 ⁱ	4,2,3
NGC 891.....	7.4	0.09	61.1	198.6	623.4	701.0	5.5	6,2,3
NGC 3627.....	10.2	0.04	56.2	144.9	507.3	458.0	4.5	4,2,3
NGC 660.....	12.0	0.08	65.5	103.8	477.4	387.0	4.2	4,2,3
NGC 3628.....	11.2	0.02	48.4	121.9	431.9	525.0	3.8	10,3
NGC 1097.....	18.0	0.10	46.7	116.1	414.0	415.0	3.6	4,2,3
NGC 3079.....	15.9	3.7	50.2	103.5	407.8	849.0	3.6	4,2,3

REFERENCES. — (1) Condon (1987); (2) Kennicutt (1998); (3) Yun, Reddy, & Condon (2001); (4) Condon et al. (1990); (5) Golla & Wielebinski (1994); (6) Condon et al. (1996); (7) Sandqvist et al. (1995); (8) Sandqvist (1999); (10) Paglione et al. (2001); (11) Condon et al. (1998)

^aDistance. In most cases, distances computed with $H = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$. In some cases, distances are taken from the literature. ^bGas surface density. Typical uncertainty in this quantity is $\pm 0.2 \text{ dex}$ (e.g., Kennicutt 1998). ^cIRAS flux density at $60 \mu\text{m}$. ^dIRAS flux density at $100 \mu\text{m}$. ^eTotal IR flux density defined by $\log_{10}(S_{\text{TIR}}) \equiv \log_{10}(2.58S_{60\mu\text{m}} + S_{100\mu\text{m}}) + \log_{10}(1.75)$, where the factor of 1.75 relates the total $[8-1000]\mu\text{m}$ flux density to the $[40-120]\mu\text{m}$ flux density (Calzetti et al. 2000; Helou et al. 1988). For any individual system listed here, this bolometric correction is uncertain at the factor of ~ 1.5 level (see Bell 2003; Dale et al. 2001). ^fRadio flux density at 1.4 GHz . ^gPredicted γ -ray flux at 1 GeV , scaled from S_{TIR} according to eq. (6), assuming $\eta = 0.05$ and $\gamma_{\text{max}} = 10^6$. ^hSee discussion of comparison with Domingo-Santamaría & Torres (2005) in §2. ⁱSee discussion of comparison with Torres (2004) in §2.

Torres, D. F., & Domingo-Santamaría, E. 2005, Modern Physics Letters A, 20, 2827
Ullio, P., Bergström, L., Edsjö, J., & Lacey, C. 2002, Phys. Rev. D, 66, 123502
Völk, H. J. 1989, A&A, 218, 67

Völk, H. J., Berezhko, E. G., & Ksenofontov, L. T. 2003, A&A, 409, 563
Yun, M. S., Reddy, N. A., & Condon, J. J. 2001, ApJ, 554, 803